

REVISITING G-DEDEKIND DOMAINS

M. ZAFRULLAH

Dedicated to fairness

ABSTRACT. Let R be an integral domain with $qf(R) = K$ and let $F(R)$ be the set of nonzero fractional ideals of R . Call R a dually compact domain (DCD) if for each $I \in F(R)$ the ideal $I_v = (I^{-1})^{-1}$ is a finite intersection of principal fractional ideals. We characterize DCDs and show that the class of DCDs properly contains various classes of integral domains, such as Noetherian, Mori and Krull domains. In addition we show that a Schreier DCD is a GCD domain with the property that for each $A \in F(R)$ the ideal A_v is principal. We show that a domain R is G-Dedekind domain (i.e. has the property that A_v is invertible for each $A \in F(R)$) if and only if R is a DCD satisfying the property $*$: for all pairs of subsets $\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\} \subseteq K \setminus \{0\}$, $(\cap_{i=1}^m (a_i))(\cap_{j=1}^n (b_j)) = \cap_{i,j=1}^{m,n} a_i b_j$. We discuss what the appropriate name for G-Dedekind domains and related notions should be. We also make some observations about how the DCDs behave under localizations and polynomial ring extensions.

1. INTRODUCTION

Let R be an integral domain with quotient field K and let $F(R)$ be the set of nonzero fractional ideals of R . Call R a dually compact domain (DCD) if for each set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ with $\cap a_\alpha R \neq (0)$ there is a finite set of elements $\{x_1, \dots, x_r\} \subseteq K \setminus \{0\}$ such that $\cap a_\alpha R = \cap_{i=1}^r x_i R$, or equivalently for each $I \in F(R)$, the ideal $I_v = (I^{-1})^{-1}$ is a finite intersection of principal fractional ideals of R . We characterize DCDs (in section 3) and show that the class of DCDs properly contains various classes of integral domains of interest such as Noetherian, Mori and Krull domains, in section 4. In addition we show that a pre-Schreier DCD is a GCD domain with the property that for each $A \in F(R)$ the ideal A_v is principal. (Here R is pre-Schreier if for all $x, y, z \in R \setminus \{0\}$ $x|yz \Rightarrow x = rs$, with $r, s \in R$ such that $r|y$ and $s|z$.) We show that a domain R is a G-Dedekind domain (i.e. has the property that A_v is invertible for each $A \in F(R)$) if and only if R is a DCD satisfying the property $*$: for all pairs of subsets $\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\} \subseteq K \setminus \{0\}$, $(\cap_{i=1}^m (a_i))(\cap_{j=1}^n (b_j)) = \cap_{i,j=1}^{m,n} a_i b_j$ from [25]. We discuss in section 2 what names for G-Dedekind domains and related notions should be appropriate. We also make some observations about how the DCDs behave under localizations and polynomial ring extensions.

1991 *Mathematics Subject Classification.* 13A15 13A05 13F05 13G05.

Key words and phrases. G-Dedekind, pseudo Dedekind, invertible, *-invertible Noetherian, Mori.

I am thankful to my friends who made me rise from the dead.

©2001 enter name of copyright holder

In [24], this author studied integral domains R with the property that A_v is invertible for every nonzero ideal A , and called these domains "generalized Dedekind Domains" or G-Dedekind domains. Later using a new form of the above mentioned \star -property, Anderson and B.G. Kang [5] published a much improved version of [24]. Calling the G-Dedekind domains "Pseudo Dedekind" domains, they showed that R is a Pseudo Dedekind domain if and only if for all sets $\{a_\alpha\}_{\alpha \in I}, \{b_\beta\}_{\beta \in J} \subseteq K \setminus \{0\}$ we have $(\cap(a_\alpha))(\cap(b_\beta)) = \cap(a_\alpha b_\beta)$, where α ranges over I and β over J . (This was actually a clever translation of the characteristic property of the G-Dedekind domains: $(AB)^{-1} = A^{-1}B^{-1}$ for all $A, B \in F(R)$ given in [24].) Call a set of elements $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ allowable if $\cap(a_\alpha) \neq (0)$. In this article, we show, among things already indicated, that for a given star operation \star , A^\star is invertible for any ideal A of R if and only if for any allowable set of elements $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ and for every nonzero ideal A we have $A^\star(\cap(a_\alpha)) = \cap A^\star a_\alpha$. As a consequence we show that R is a Pseudo Dedekind/G-Dedekind domain if and only if for each nonzero ideal A of R and for each allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $A_v(\cap(a_\alpha)) = \cap A_v(a_\alpha)$. We show that R is a DCD if and only if for any allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ there is a set $\{b_1, b_2, \dots, b_n\} \subseteq K \setminus \{0\}$ such that $\cap_{\alpha \in I}(a_\alpha) = (b_1, b_2, \dots, b_n)_v$. We also show that R is a G-Dedekind/Pseudo Dedekind domain if and only if R is a DCD with the above mentioned property \star . That is, the DCDs were essentially at work behind the scenes in the results in [24] and subsequently in [5].

We use star operations, in order to approach the subject from a more general standpoint. Included below is a brief introduction to star operations. The reader may consult [15], [11] or [16] for more information on star operations. For our purposes we include below some information that may be helpful in reading this article.

Let R be an integral domain with quotient field K and let $F(R)$ be the set of nonzero fractional ideals of R . A star operation is a function $A \mapsto A^\star$ on $F(R)$ with the following properties:

- If $A, B \in F(R)$ and $a \in K \setminus \{0\}$, then
- (i) $(a)^\star = (a)$ and $(aA)^\star = aA^\star$.
- (ii) $A \subseteq A^\star$ and if $A \subseteq B$, then $A^\star \subseteq B^\star$.
- (iii) $(A^\star)^\star = A^\star$.

We may call A^\star the \star -image (or \star -envelope) of A . An ideal A is said to be a \star -ideal if $A^\star = A$. Thus A^\star is a \star -ideal (by (iii)). Moreover (by (i)) every principal fractional ideal, including $R = (1)$, is a \star -ideal for any star operation \star .

For all $A, B \in F(R)$ and for each star operation \star , we can show that $(AB)^\star = (A^\star B)^\star = (A^\star B^\star)^\star$. These equations define what is called \star -multiplication (or \star -product). Associated with each star operation \star is a star operation \star_f defined by $A^{\star_f} = \bigcup \{J^\star \mid 0 \neq J \text{ is a finitely generated subideal of } A\}$, for each $A \in F(R)$. We say that a star operation \star is of finite type or of finite character if $\star = \star_f$.

Define $A^{-1} = \{x \in K \mid xA \subseteq R\}$. Thus $A^{-1} = \cap_{a \in A \setminus \{0\}} (\frac{1}{a})$. Also define $A_v = (A^{-1})^{-1}$ and $A_t = A_{v_f} = \bigcup \{J_v \mid 0 \neq J \text{ is a finitely generated subideal of } A\}$. By the definition $A_t = A_v$ for each finitely generated nonzero ideal of R . The functions $A \mapsto A_v$ and $A \mapsto A_t$ on $F(R)$ are more familiar examples of star operations defined on an integral domain. A fractional ideal $A \in F(R)$ is \star -invertible if $(AA^{-1})^\star = R$. An invertible ideal is a \star -invertible \star -ideal for each \star -operation \star and so is a v -ideal. A v -ideal is better known as a divisorial ideal and using the definition it can be shown that $A_v = \bigcap_{\substack{x \in K \setminus \{0\} \\ A \subseteq xR}} xR$. The identity function d on $F(R)$, defined

by $A \mapsto A$ is another example of a star operation. Indeed a " d -invertible" ideal is the usual invertible ideal. There are of course many more star operations that can be defined on an integral domain R . But for any star operation \star and for any $A \in F(R)$, $A^\star \subseteq A_v$. Some other useful relations are: For any $A \in F(R)$, $(A^{-1})^\star = A^{-1} = (A^\star)^{-1}$ and so, $(A_v)^\star = A_v = (A^\star)_v$. Using the definition of the t -operation one can show that an ideal that is maximal w.r.t. being a proper integral t -ideal is a prime ideal of R , each nonzero ideal A of R with $A_t \neq R$ is contained in a maximal t -ideal of R and $R = \cap R_M$, where M ranges over maximal t -ideals of R . The set of maximal t -ideals of R is denoted by $t\text{-Max}(R)$. For more on v - and t -operations the reader may consult sections 32 and 34 of Gilmer [15], or the other two books cited above. This is the barest minimum of description to get us started, we shall expand on it when need arises. Our terminology comes from [15]. Of course we have called a subset $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ allowable $\cap(a_\alpha) \neq (0)$ (a) to save on space and (b) because the characteristic property of a G-Dedekind/pseudo Dedekind domain is that for all $A, B \in F(D)$ we have $(AB)^{-1} = A^{-1}B^{-1}$ and this does not require any set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ with $\cap(a_\alpha) = (0)$. The reader may expect such purpose oriented terminology in the sequel as well. In section 2 we settle a name for G-/Pseudo Dedekind domains, indicating reasons why we should, and in section 3 we show that R is a G-/Pseudo Dedekind domain if and only if R is a DC \star -domain. Finally in section 4 we touch on some related questions and indicate how DCDs behave under some extensions such as quotient ring formation or, in case of integrally closed DCDs, polynomial ring formation.

2. WHAT'S IN A NAME?

Popescu [20] introduced the notion of a generalized Dedekind domain via localizing systems. Nowadays, the following equivalent definition is usually given: an integral domain is a generalized Dedekind domain if it is a strongly discrete Prüfer domain (i.e., $P \neq P^2$ for every prime ideal P) and every (prime) ideal I has $\sqrt{I} = \sqrt{(a_1, \dots, a_n)}$ for some $a_1, \dots, a_n \in I$ (or equivalently, every principal ideal has only finitely many minimal prime ideals). Unbeknownst to this author [20] was already out. I personally do not think there is anything pseudo about the G-Dedekind domains. On the other hand some serious studies related to G-Dedekind prime rings, introduced by Evrim Akalan [9], are being carried out. Hence the need for a name close to G-Dedekind domains. The aim of this section is to fix a suitable name for the domains that are given two different names, one not quite appropriate and the other overshadowed by a previously adopted name. Of course as a result we end up with one more notion a more suitable name for a so called π -domain.

The following lemma essentially comes from [1], yet we use it to give a more general view of what started as G-Dedekind domains or Pseudo Dedekind domains. Of course our statement is different and more streamlined.

Lemma 2.1. *Let \star be a star operation defined on an integral domain R and let $A \in F(R)$. Then A^\star is invertible if and only if for any allowable set of elements $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $(A^\star(\cap(a_\alpha))) = \cap A^\star a_\alpha$.*

The proof of the above lemma has been used as part of the proof of the following theorem.

Theorem 2.2. *Let \star be a star operation defined on an integral domain R . Then A^\star is invertible for every nonzero fractional ideal A of R if and only if for any allowable set of elements $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ and for every nonzero ideal A we have $(A^\star(\cap(a_\alpha))) = \cap A^\star a_\alpha$.*

Proof. Let A be any nonzero ideal of R and suppose that for every allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $A^\star(\cap(a_\alpha)) = \cap A^\star a_\alpha$. Then $R \supseteq A^\star A^{-1} = A^\star(\cap_{a_\beta \in A \setminus \{0\}}(\frac{1}{a_\beta})) = (\cap_{a_\beta \in A \setminus \{0\}} A^\star(\frac{1}{a_\beta}))$ by the condition. Since for each of $a_\beta \in A \setminus \{0\}$ we have $A^\star(\frac{1}{a_\beta}) \supseteq R$ and so $R \supseteq (A^\star A^{-1} = A^\star(\cap_{a_\beta \in A \setminus \{0\}}(\frac{1}{a_\beta})) = (\cap_{a_\beta \in A \setminus \{0\}} A^\star(\frac{1}{a_\beta}))) \supseteq R$, showing that $R = A^\star(\cap_{a_\beta \in A \setminus \{0\}}(\frac{1}{a_\beta})) = A^\star A^{-1}$. Thus, as A was chosen arbitrarily, the condition implies that for every nonzero ideal A we have that A^\star is invertible. Conversely suppose that A is a nonzero ideal such that A^\star is invertible. Then, by an exercise on page 80 of [15] we have, for any allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, $A^\star(\cap(a_\alpha)) = \cap A^\star(a_\alpha)$. Thus for every nonzero ideal A the ideal A^\star being invertible implies that for every nonzero ideal A , and for every allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $A^\star(\cap(a_\alpha)) = \cap A^\star a_\alpha$. \square

Corollary 1. *For $\star = d$, R is a Dedekind domain if and only if for each nonzero ideal A of R and for each allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $A(\cap(a_\alpha)) = \cap A(a_\alpha)$.*

Proof. Indeed it is well known that R is a Dedekind domain if and only if every nonzero ideal of R is invertible and Theorem 2.2 provides the, general, necessary and sufficient conditions for every nonzero ideal to be \star -invertible, when $\star = d$. \square

Let's recall that an ideal I is t -invertible if I is v -invertible and I^{-1} is a v -ideal of finite type, that an ideal I that is invertible, is \star -invertible for every star operation \star [27] and that an integral domain R is a Krull domain if and only if every nonzero ideal of R is t -invertible [19]. Now if I_t is invertible, then I_t and hence I is t -invertible. Thus if, for each ideal I , I_t is invertible in a domain R then, R is at least a Krull domain. According to Theorem 1.10 of [24] if I_t is invertible for every nonzero ideal of R , then R is a locally factorial Krull domain. Such domains are often called π -domains for some reason. Now Theorem 2.2 characterizes π -domains for $\star = t$ as follows.

Corollary 2. *A domain R is a π -domain if and only if for each nonzero ideal A of R and for each allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $A_t(\cap(a_\alpha)) = \cap A_t(a_\alpha)$.*

Proof. Indeed R is a π -domain if and only if A_t is invertible for every nonzero ideal A of R (Theorem 1.10 of [24]) and Theorem 2.2 provides the, general, necessary and sufficient conditions for every nonzero ideal to be \star -invertible, when $\star = t$. \square

Of course by saying that an ideal A of R is v -invertible we mean that $(AA^{-1})_v = R$. Similar to earlier comments we note that A_v being invertible entails A being v -invertible. We also note that the domains R with A_v invertible for each nonzero ideal A are the G -Dedekind/Pseudo Dedekind domains. So for $\star = v$, Theorem 2.2 provides the following characterization of G -Dedekind/Pseudo Dedekind domains.

Corollary 3. *A domain R is a G -Dedekind/Pseudo Dedekind domain if and only if for each nonzero ideal A of R and for each allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, we have $A_v(\cap(a_\alpha)) = \cap A_v(a_\alpha)$.*

The proof is the same as the one provided by Theorem 2.2 replacing \star by v .

Remark 2.3. Looking at Theorem 2.2 and Corollaries 1, 2, 3, we may call R a \star -Dedekind domain if A^\star is invertible for each $A \in F(R)$ and note that in a \star -Dedekind domain we have $A^\star = A_v$ for all $A \in F(R)$. This is because an invertible ideal is divisorial. Thus Corollary 2 gives for $\star = t$ a t -Dedekind domain and Corollary 3 gives the name of a v -Dedekind domain to the G-Dedekind domain of [24] and Pseudo Dedekind domain of [5]. But there is a slight problem with this naming system, Jesse Elliott in [11] calls a Krull domain a t -Dedekind domain. So, perhaps, \star -G-Dedekind may be the general name with the note that a d -G-Dedekind domain is the usual Dedekind domain and a t -G-Dedekind domain is a locally factorial Krull domain while the v -G-Dedekind domain is the usual Pseudo Dedekind domain, or the old G-Dedekind domain. Of course, as $\star = v$ in a \star -G-Dedekind domain, each \star -G-Dedekind domain has the properties listed in [24] for G-Dedekind domains are shared by \star -G-Dedekind domains, or in [5] for Pseudo Dedekind domains. Thus if \star is of finite character, then $\star = \star_f = v_f = t$. That is if R is a \star -G-Dedekind domain and \star is of finite character, then R is a t -G-Dedekind domain. Recall that an integral domain R with quotient field K is completely integrally closed if whenever $rx^n \in R$ for $x \in K$, $0 \neq r \in R$, and every integer $n \geq 1$, then $x \in R$. Equivalently, R is completely integrally closed if and only if $(AA^{-1})_v = R$ for every $A \in F(R)$ [15]. If, for a star operation \star defined on R , A is \star invertible for each $A \in F(R)$ following [3] we may call R a \star -CICD. Thus, as noted in [24], a \star -G-Dedekind domain is a completely integrally closed domain (CICD), for each star operation \star that it is defined for. There is a star operation called the w -operation, defined in terms of the t -operation as $A \mapsto A_w = \bigcap_{M \in t\text{-Max}(R)} AR_M$, see [4] and references there. As indicated in [4], this operation is of finite character. Thus, in view of earlier comments in this remark, a w -G-Dedekind domain is a t -G-Dedekind domain.

3. DUALY COMPACT DOMAINS

Cohn [10] called an element $x \in R$ primal if for all $y, z \in R$, $x|yz$ implies that $x = rs$ where $r|y$ and $s|z$. A domain all of whose nonzero elements are primal was called a pre-Schreier domain in [25]; this was a break from Cohn who called R a Schreier domain if R was integrally closed with all elements primal. Based on a study of the group of divisibility of a pre-Schreier domain this author extracted [25] what he called the \star property: R is a \star domain if for all pairs of subsets $\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\} \subseteq K \setminus \{0\}$, $(\bigcap_{i=1}^m (a_i))(\bigcap_{j=1}^n (b_j)) = \bigcap_{i,j=1}^{m,n} a_i b_j$.

We now look at the facts working behind the Anderson-Kang/Zafrullah results. For this let us call a domain R dually compact (DC) if for any allowable subset $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, there is a set $\{a_1, a_2, \dots, a_n\} \subseteq K \setminus \{0\}$ such that $\bigcap_{\alpha \in I} (a_\alpha) = (a_1) \cap (a_2) \cap \dots \cap (a_n)$. Let's also note that a fractional ideal A being divisorial ideal (or v -ideal) of finite type means that there are elements $x_1, \dots, x_r \in K \setminus \{0\}$ such that $A = (x_1, \dots, x_r)_v$.

Theorem 3.1. *The following are equivalent for an integral domain R that is different from its quotient field K .*

- (1) R is DC,
- (2) A_v is a v -ideal of finite type for every $A \in F(R)$,
- (3) A^{-1} is of finite type for each $A \in F(R)$,

- (4) A^{-1} is of finite type for each nonzero integral ideal A of R ,
- (5) A_v is of finite type for each nonzero integral ideal A of R ,
- (6) every divisorial ideal of R is expressible as a finite intersection of principal fractional ideals,
- (7) for a star operation \star and for each $A \in F(R)$, the ideal A^\star is a v -ideal of finite type,
- (8) for a star operation \star and for each $A \in F(R)$, the ideal A^\star is a finite intersection of principal ideals from $F(R)$.

Proof. (1) \Rightarrow (2). Suppose that R is DC and suppose that A is a nonzero fractional ideal generated by $\{c_\gamma\}_{\gamma \in J}$. Then $A^{-1} = \cap_\gamma (\frac{1}{c_\gamma}) \neq (0)$. So by the DC property there is a finite set $\{x_1, x_2, \dots, x_n\} \subseteq K \setminus \{0\}$ such that $\cap_\gamma (\frac{1}{c_\gamma}) = \cap_{i=1}^n (x_i) = (x_1^{-1}, \dots, x_n^{-1})^{-1}$. Now $A^{-1} = (x_1^{-1}, \dots, x_n^{-1})^{-1}$ implies that $A_v = (x_1^{-1}, \dots, x_n^{-1})_v$.

(2) \Rightarrow (1). Suppose that for each $A \in F(R)$, there are $x_1, x_2, \dots, x_n \in K \setminus \{0\}$ such that $A_v = (x_1, x_2, \dots, x_n)_v$. Then since A^{-1} is a divisorial ideal, as $(A^{-1})_v = A^{-1}$, we have $A^{-1} = (y_1, \dots, y_r)_v$. Or, assuming that all the y_i are nonzero, $A_v = \cap (\frac{1}{y_i})$ for each $A \in F(R)$. Thus if for each $A \in F(R)$ A_v is of finite type, then for each $A \in F(R)$ we can find some $b_i \in K \setminus \{0\}$ such that $A_v = \cap_{i=1}^r (b_i)$. Now let $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ be allowable and let $A = \cap (a_\alpha)$. Since $\cap (a_\alpha) \neq (0)$, A is a divisorial ideal by [15] and hence of finite type and so, by (2) for some $x_1, \dots, x_n \in K \setminus \{0\}$ we have $A = \cap (a_\alpha) = \cap_{i=1}^n (x_i)$.

Next (2) \Leftrightarrow (5) and (3) \Leftrightarrow (4) because every fractional ideal A of R is of the form $\frac{B}{d}$ where B is an integral ideal. (2) \Rightarrow (3) because $A^{-1} = (A^{-1})_v$ and (3) \Rightarrow (2) because $A_v = (A^{-1})^{-1}$.

(1) \Rightarrow (6). A nonzero ideal A is divisorial if and only if $A = \cap_{\substack{A \subseteq x_\alpha R \\ x \in K \setminus \{0\}}} x_\alpha R$. By the DC condition there are x_1, \dots, x_n in $K \setminus \{0\}$ such that $A = \cap_{\substack{A \subseteq x_\alpha R \\ x \in K \setminus \{0\}}} x_\alpha R = \cap_{i=1}^n x_i R$.

(6) \Rightarrow (1). Let for $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, $\cap (a_\alpha) \neq (0)$. Note that $\cap (a_\alpha)$ is divisorial. So by (6) there are elements $a_1, \dots, a_r \in K \setminus \{0\}$ such that $\cap (a_\alpha) = \cap_{i=1}^r (a_i)$. Finally each of (7) and (8) holds if and only if $A^\star = A_v$ for all $A \in F(R)$ because in both cases $A^\star = B$ where B is divisorial and so $A_v = B = A^\star$. Observing that, we have (7) \Leftrightarrow (5) and (8) \Leftrightarrow (6). \square

Remark 3.2. (1) There are a number of integral domains that fit the description of DCDs. Noetherian domains do fit nicely, as do the so-called Mori domains. Recall that R is a Mori domain if R satisfies ascending chain conditions on divisorial ideals. It is well known that R is a Mori domain if and only if for each $A \in F(R)$ there is a finitely generated fractional ideal $B \subseteq A$ such that $A_v = B_v$. Indeed a Krull domain is a DCD, being a Mori domain as indicated in Fossum [12]. On the other hand, there are DCDs, such as the ring of entire functions in which A_v is principal and, the ring of entire functions is neither a Mori domain, nor a Krull domain. (2) If the star operation \star defined on R is such that A^\star is a v -ideal of finite type for each $A \in F(R)$ we can call R a \star -DCD. (3) It may be somewhat hard to see, for some, that any t -invertible t -ideal, and hence any invertible ideal, is a finite intersection of principal fractional ideals. Let me note for the record that if A is a t -invertible t -ideal, then A^{-1} is of finite type. Say for some $B = \{b_1, \dots, b_n\}$ we have $A^{-1} = B_v$. But then $A = A_v = (B_v)^{-1} = \cap_{i=1}^n (\frac{1}{b_i})$.

Indeed as in a DCD the inverse of every nonzero fractional ideal is of finite type, all we need for the v -G-Dedekind property to hold is the $*$ -property.

Theorem 3.3. *Let R be a DCD. Then R is a v -G-Dedekind domain if and only if R is a $*$ -domain.*

Proof. It is easy to see, from the treatment of it in [5], that a v -G-Dedekind domain is a $*$ -domain. (This fact was also mentioned in [24].) But of course we need to show that a v -G-Dedekind domain is DC. This follows from the fact that, in a v -G-Dedekind domain R , A_v is invertible and, as shown in Remark 3.2, an intersection of finitely many principal fractional ideals. For the converse we show that DC plus the $*$ -property implies the v -G-Dedekind property. For this consider for allowable sets $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$, $\{b_\beta\}_{\beta \in J} \subseteq K \setminus \{0\}$ the product $P = (\cap(a_\alpha))(\cap(b_\beta))$. By the DC property of R we can find $\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\} \subseteq K \setminus \{0\}$ such that $\cap(a_\alpha) = \cap_{i=1}^m(a_i)$ and $\cap(b_\beta) = (\cap_{j=1}^n(b_j))$. Thus using DC plus $*$, $P = (\cap(a_\alpha))(\cap(b_\beta)) = (\cap(a_i))(\cap(b_j)) = \cap_{i,j=1}^{m,n} a_i b_j = \cap_i a_i ((\cap(b_j)) = \cap a_i (\cap(b_j))$ (using $\cap(b_j) = \cap(b_\beta)$). This gives $P = \cap_i a_i (\cap(b_\beta)) = \cap_{i,\beta} a_i b_\beta = \cap_\beta b_\beta (\cap(a_i)) = \cap_\beta b_\beta (\cap(a_\alpha)) = \cap_{\alpha,\beta} (a_\alpha b_\beta)$. \square

Recall that an integral domain R is called v -coherent if every finite intersection of v -ideals of finite type is a v -ideal of finite type, equivalently, if A^{-1} is a v -ideal of finite type for all finitely generated $A \in F(R)$, [13]. Also that R is a GGCD domain if $aR \cap bR$ is invertible for all $a, b \in R \setminus \{0\}$, [2]. According to Corollary 1.7 of [24] a v -coherent domain R is a GGCD domain if and only if R is a $*$ -domain. Now a DCD R is slightly more than a v -coherent domain.

Corollary 4. *Suppose that for a star operation \star , A^\star is a v -ideal of finite type for each $A \in F(R)$. Then R is a \star -G-Dedekind domain if and only if R is a $*$ -domain. Consequently a Mori (and hence a Krull) domain R is a locally factorial Krull domain if and only if R is a $*$ -domain.*

The proof can be easily constructed from the preceding comments and so is left to the reader.

Considered in [5] was also the notion of a pseudo principal ideal domain (pseudo PID) or, in our terminology, v -G-PID by requiring that A_v is principal, for each nonzero ideal A of R . This notion appeared in Bourbaki too, as pointed out in [5]. Using the DC approach, one can prove the following result.

Theorem 3.4. *A domain R is a v -G-PID if and only if R is a DC Schreier domain.*

Proof. Suppose that R is a DC Schreier domain. Now a Schreier domain is a $*$ -domain too [25, Corollary 1.7]. So a DCD that is a Schreier domain is at least a v -G-Dedekind domain, by Theorem 3.3. But then A_v is invertible for each nonzero ideal A of R and in a Schreier domain every invertible ideal is principal [25, Theorem 3.6]. Thus A_v is principal for each nonzero ideal A of R and R is a v -G-PID. Conversely note that a v -G-PID is DC and is at least a GCD domain and a GCD domain is Schreier [10]. Thus a v -G-PID R is a DC Schreier domain. \square

Theorem 3.5. *Let \star be a star operation defined on R such that for each $A \in F(R)$, A^\star is a v -ideal of finite type. Then the following are equivalent.*

- (1) R is a \star -G-PID,
- (2) R is a Schreier domain,
- (3) R is a \star -G-Dedekind domain with $\text{Pic}(R) = (0)$.

Proof. (1) \Leftrightarrow (2). A \star -G-PID, by definition, is a GCD domain and so a Schreier domain. Conversely, "for each $A \in F(R)$, A^* is a v -ideal of finite type" makes each $A^* \in F(R)$ a v -ideal of finite type. But then for each $A \in F(D)$ A^* is a finite intersection of principal fractional ideals and of finite type and by Theorem 3.6 of [25] principal because R is Schreier.

(1) \Leftrightarrow (3). \star -GPID is \star -G-Dedekind with every invertible ideal principal which is exactly (3).

For the converse note that (3) implies that R is at least a GCD domain and a GCD domain is Schreier. That is (2) holds and (2) is equivalent to (1). \square

4. RELATED STUFF

We end this article with some interesting characterizations of the \star -GPIDs and discussion of related material.

Recall that a Riesz group is a directed group that satisfies the Riesz interpolation property: given that $x_1, x_2, \dots, x_m; y_1, y_2, \dots, y_n \in G$ such that $x_i \leq y_j$ for all $i \in [1, m], j \in [1, n]$ there is $z \in G$ such that $x_i \leq z \leq y_j$ for all $(i, j) \in [1, m] \times [1, n]$. It was shown in [25] that the Riesz interpolation property translates in the commutative ring theory set up to: for all $x_1, x_2, \dots, x_m; y_1, y_2, \dots, y_n \in K \setminus \{0\}$ with $x_1, x_2, \dots, x_m \in \cap y_i R$ there is a $z \in \cap y_i R$ such that $(x_1, \dots, x_m) \subseteq zR \subseteq \cap y_i R$. So a pre-Schreier domain is actually a pre-Riesz domain. We have no interest in changing existing names, we only want to add a new name. Call R a super Riesz domain if for any divisorial ideal A of R and for any set $\{x_\alpha\}$ of elements contained in A , with $\cap(x_\alpha) \neq (0)$, there is a $d \in A$ such that $(x_\alpha) \subseteq (d)$. Because a divisorial ideal is expressible as an intersection of principal fractional ideals, a super Riesz domain can be easily seen to be pre-Schreier. Also let's call a product AB of ideals A, B subtle if for each $x \in AB$ we have $x = ab$ where $a \in A$ and $b \in B$. The authors of [5] also touched on the following question: Let R be an integral domain that satisfies $(\cap(a_\alpha))(\cap(b_\beta)) = \cap(a_\alpha b_\beta)$ for all subsets $\{a_\alpha\}_{\alpha \in I} \subseteq R \setminus \{0\}, \{b_\beta\}_{\beta \in J} \subseteq R \setminus \{0\}$. Is R pseudo-Dedekind?

We try to give a partial answer below.

Proposition 1. *Let R be an integral domain. Then the following are equivalent.*

(1) For all allowable $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}, \{b_\beta\}_{\beta \in J} \subseteq K \setminus \{0\}$ we have $(\cap(a_\alpha))(\cap(b_\beta)) = \cap(a_\alpha b_\beta)$,

(2) for all allowable $\{a_\alpha\}_{\alpha \in I} \subseteq R \setminus \{0\}, \{b_\beta\}_{\beta \in J} \subseteq R \setminus \{0\}$ we have $(\cap(a_\alpha))(\cap(b_\beta)) = \cap(a_\alpha b_\beta)$ and for all allowable sets $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ we have a $d \in R \setminus \{0\}$ such that $d(\cap(a_\alpha))$ is expressible as an intersection of principal integral ideals.

(3) R is a DCD that satisfies: for all allowable $\{a_\alpha\}_{\alpha \in I} \subseteq R \setminus \{0\}, \{b_\beta\}_{\beta \in J} \subseteq R \setminus \{0\}$ we have $(\cap(a_\alpha))(\cap(b_\beta)) = \cap(a_\alpha b_\beta)$.

(1) \Rightarrow (2). Obvious because the first part follows directly and (1) means R is DC and so for each allowable set $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}$ we can find a_1, \dots, a_n such that $(\cap(a_\alpha)) = (\cap_{i=1}^n(a_i))$. But for the right hand we can find a nonzero d such that da_i are all integral.

(2) \Rightarrow (1). Consider $(\cap(a_\alpha))(\cap(b_\beta))$ for all allowable $\{a_\alpha\}_{\alpha \in I} \subseteq K \setminus \{0\}, \{b_\beta\}_{\beta \in J} \subseteq K \setminus \{0\}$. Since $(\cap(a_\alpha)), (\cap(b_\beta))$ are nonzero fractional ideals by the given property of R we have, for some $r, s \in R$, $r(\cap(a_\alpha)) = \cap(x_\gamma)$ and $s(\cap(b_\beta)) = \cap(y_\delta)$, where x_γ and y_δ are in R , by (2). But then $(\cap(x_\gamma))(\cap(y_\delta)) = \cap(x_\gamma y_\delta) = P$. Now $P = \cap(x_\gamma y_\delta) = \cap_\gamma x_\gamma ((\cap(y_\delta)))$ (substituting for $\cap(y_\delta) = \cap_\gamma x_\gamma ((\cap(y_\delta))) = \cap_{\gamma, \beta} (x_\gamma s b_\beta) =$

$\cap_{\beta} s b_{\beta} \cap (x_{\gamma})$ and substituting for $\cap(x_{\gamma})$ we get $P = \cap_{\beta} s b_{\beta} \cap (r a_{\alpha}) = \cap (r a_{\alpha} s b_{\beta}) = r s (\cap (a_{\alpha} b_{\beta}))$. But on the other hand $P = (\cap(x_{\gamma})) (\cap(y_{\delta})) = (r (\cap(a_{\alpha}))) (s (\cap(b_{\beta})))$. Thus $r s ((\cap(a_{\alpha}))) ((\cap(b_{\beta}))) = r s (\cap(a_{\alpha} b_{\beta}))$. Canceling $r s$ from both sides we get the desired equality.

(1) \Rightarrow (3). Obvious, in light of (1) \Rightarrow (2).

(3) \Rightarrow (2). Note that because R is DC for each allowable set $\{a_{\alpha}\}_{\alpha \in I} \subseteq K \setminus \{0\}$ we can find a_1, \dots, a_n such that $(\cap(a_{\alpha})) = (\cap_{i=1}^n (a_i))$ and so a $d \in R \setminus \{0\}$ such that $d a_i \in R$. But then $d(\cap(a_{\alpha})) = \cap_{i=1}^n d a_i$ an intersection of principal fractional ideals.

Recall from [25], again, that R is a pre-Schreier domain if and only if for all $\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\} \subseteq R \setminus \{0\}$, $a_i b_j | x$ implies $x = r s$ where $a_i | r$ and $b_j | s$. Since this property sprang in the context of pre-Schreier domains we can call a domain R super pre-Schreier if $\{a_{\alpha}\}_{\alpha \in I} \subseteq R \setminus \{0\}$, $\{b_{\beta}\}_{\beta \in J} \subseteq R \setminus \{0\}$ such that $a_{\alpha} b_{\beta} | x$ then $x = r s$ such that $a_{\alpha} | r$ and $b_{\beta} | s$, and ask if a super pre-Schreier domain must be a v -GPID. The reason for this question is provided by the following proposition.

Proposition 2. *An integral domain R is super pre-Schreier if and only if for all allowable $\{a_{\alpha}\}_{\alpha \in I} \subseteq R \setminus \{0\}$, $\{b_{\beta}\}_{\beta \in J} \subseteq R \setminus \{0\}$ we have $(\cap(a_{\alpha}))(\cap(b_{\beta})) = (\cap(a_{\alpha} b_{\beta}))$ such that for each $x \in (\cap(a_{\alpha}))(\cap(b_{\beta}))$ $x = r s$ where $r \in (\cap(a_{\alpha}))$ and $s \in (\cap(b_{\beta}))$.*

Proof. Suppose that R is super pre-Schreier. Then $(\cap(a_{\alpha}))(\cap(b_{\beta})) \subseteq (\cap(a_{\alpha} b_{\beta}))$ always holds. So let $x \in (\cap(a_{\alpha} b_{\beta}))$. This means $a_{\alpha} b_{\beta} | x$. But super pre-Schreier property requires that $x = r s$ where $r \in \cap(a_{\alpha})$ and $s \in \cap(b_{\beta})$ and that puts $x \in (\cap(a_{\alpha}))(\cap(b_{\beta}))$ and the other requirement is met. The converse is a similar translation. \square

Call the product IJ of two nonzero integral ideals I, J of R subtle if each $d \in IJ \setminus \{0\}$ can be written as $d = r s$ where $r \in I$ and $s \in J$. It was shown in [25] that R is pre-Schreier if and only if R has the property $*$ and for each pair of subsets $\{a_1, \dots, a_m\}, \{b_1, \dots, b_n\} \subseteq R \setminus \{0\}$ the product $(\cap_{i=1}^m (a_i)) (\cap_{i=1}^n (b_i))$ is subtle. (Following an earlier version of [25], Anderson and Dobbs [7] studied domains products of whose ideals were all subtle.)

Corollary 5. *The following are equivalent for an integral domain R .*

(1) For all allowable $\{a_{\alpha}\}_{\alpha \in I} \subseteq R \setminus \{0\}$, $\{b_{\beta}\}_{\beta \in J} \subseteq R \setminus \{0\}$ we have $(\cap(a_{\alpha}))(\cap(b_{\beta})) = (\cap(a_{\alpha} b_{\beta}))$ such that for each $x \in (\cap(a_{\alpha}))(\cap(b_{\beta}))$ $x = r s$ where $r \in (\cap(a_{\alpha}))$ and $s \in (\cap(b_{\beta}))$,

(2) for every pair of nonzero ideals A, B we have $(AB)_v = A_v B_v$, for every nonzero integral ideal A of R there is a $d \in R \setminus \{0\}$ with $d A_v$ an intersection of principal integral ideals and the product is subtle,

(3) R is a super pre-Schreier domain.

Finally, a word about DCDs. Indeed a DCD can be characterized by: A_v is a v -ideal of finite type for each $A \in F(R)$. The first thing that comes to mind as a general property is that A^{-1} is of finite type for each $A \in F(R)$. This leads to the following result. But we need to recall some terminology. R is \star -Prüfer if for each finitely generated $A \in F(R)$ is \star -invertible [3]. Since, for $A \in F(R)$, A being \star -invertible implies $A^{\star} = A_v$. So for each finitely generated nonzero ideal J in a \star -Prüfer domain R we have $J^{\star} = J_v$. Also as $(JJ^{-1})^{\star} = R$ implies $((JJ^{-1})^{\star})_v = (JJ^{-1})_v = R$, a \star -Prüfer domain is a v -domain and $\star_f = t$. If \star is of finite character, then A^{-1} is a v -ideal of finite type for each finitely generated $A \in F(R)$ and so, if \star is of finite type, a \star -Prüfer domain is a Prüfer \star -multiplication domain and

$\star = t$. \star -Prüfer domains, for a finite character \star -operation \star , were studied in [18] where they were called \star -multiplication domains. They were later called Prüfer \star -multiplication domains (P \star MDs). An in-depth study of these domains can be found in [14], along with an introduction to semistar operations. These days this concept is defined by: R is a P \star MDs, for a star operation \star , if for each finitely generated $A \in F(R)$, A is \star_f -invertible. If R is a P \star MD for a finite character star operation, then $\star = t$ over R . On the other hand if R is a P \star MD for a "general" star-operation \star , then $\star_f = t$.

Proposition 3. *Let R be a DCD and let \star be a star operation defined on R . If A is \star -invertible, for $A \in F(R)$, then A^\star is \star_f -invertible. Consequently a DCD R is a P \star MD if and only if R is a \star -Prüfer domain.*

Proof. Note that A being \star_f -invertible means that A^{-1} is a v -ideal of finite type. But A^{-1} is a v -ideal of finite type in a DCD. Thus a DC \star -Prüfer domain is a P \star MD. The converse is obvious in the case of a DCD. \square

Remark 4.1. It may be noted that there is a marked difference between " A is \star_f -invertible" and " A^\star is \star_f -invertible". That is if you require that every $A \in F(R)$ is \star_f -invertible, you will end up with a Krull domain, for \star_f -invertible is t -invertible and every $A \in F(R)$ being t -invertible implies that R is Krull [19, page 82]. On the other hand if you require that for each $A \in F(R)$ the ideal A^\star is \star_f -invertible, you get a PVMD that acts and behaves very much like \star -G-Dedekind domains. These domains were studied under the name of pre-Krull domains in [26], just for $\star = v$ and later, under the name of (t, v) -Dedekind domains, in [3], essentially based on the line taken in [26], describing the (t, v) -Dedekind domains as domains in which $(AB)^{-1} = (A^{-1}B^{-1})^t$, or as domains in which A_v is t -invertible for each $A \in F(R)$. The closest to that [3] came to was in its Theorem 1.14. These and similar concepts were also studied by Halter-Koch under "mixed invertibility". Hopefully, with the introduction of DC property the situation will become clearer.

For now we have the following corollary.

Corollary 6. *The following are equivalent for an integral domain R , with a star operation \star defined on it.*

- (1) A^\star is \star_f -invertible for each $A \in F(R)$,
- (2) In R we have, A \star -invertible and $(AB)^\star = (A^\star B^\star)^{\star_f}$, for all $A, B \in F(R)$,
- (3) R is a \star -DC \star -Prüfer domain.
- (4) R is a \star -CICD and $(AB)^\star = (A^\star B^\star)^{\star_f}$, for all $A, B \in F(R)$.

Proof. (1) \Rightarrow (2). Let A^\star be \star_f -invertible, for all $A \in F(R)$. Then A^\star and hence A is \star -invertible, forcing $A^\star = A_v$ [27], for all $A \in F(R)$. Thus $\star = v$ over R . Moreover, as \star_f is of finite character, A^\star is a \star -ideal of finite type, because A^\star is \star_f -invertible. Now consider for $A, B \in R$ the product $(AB)^\star$. Indeed $(AB)^\star = (A^\star B^\star)^\star$. Because each of A^\star, B^\star is of finite type $(AB)^\star = (A^\star B^\star)^{\star_f}$.

(2) \Rightarrow (1). $R = (AA^{-1})^\star = (A^\star A^{-1})^{\star_f}$, by (2).

(2) \Rightarrow (3). By (2) A is \star -invertible for every $A \in F(R)$ we conclude that R is a \star -CICD and hence \star -Prüfer. Since by (2) we have $(AB)^\star = (A^\star B^\star)^{\star_f}$ we conclude that $R = (AA^{-1})^\star = (A^\star A^{-1})^{\star_f}$ which established that A^\star is a \star -ideal of finite type. But as $\star = v$ we conclude that R is \star -DC.

(3) \Rightarrow (1). Follows from Proposition 3.

Finally, (4) and (2) are re-statements of each other. \square

The domain characterized by Corollary 6 is a completely integrally closed PVMD, that is a Dedekind domain for $\star = d$ a Krull domain for \star of finite character, or for $\star = t$, and a pre-Krull domain of [26] for a star operation \star that is not of finite type, or $\star = v$. The following proposition is a reason why I tend to call these domains (\star, v) -G-Dedekind domains. For this let us recall that R is \star -DC if for each $A \in F(R)$ we have that A^\star is a v -ideal of finite type.

Proposition 4. *A (\star, v) -G-Dedekind domain R is a \star -G-Dedekind domain if and only if R is a \star -domain.*

Proof. A (\star, v) -G-Dedekind domain R is \star -DC, with $\star = v$, and so DC while a DC \star -domain is indeed a \star -G-Dedekind domain, by Theorem 3.3. Conversely a \star -G-Dedekind domain is a \star -domain and DC. This means that it is \star -DC and as every \star -ideal in a \star -G-Dedekind domain is invertible we conclude that a \star -G-Dedekind domain is a \star -domain that is a (\star, v) -G-Dedekind domain for every finite type star operation \star . \square

It may be noted that while (\star, v) -G-Dedekind domains have been studied before, Corollary 6 and Proposition 4 provide a better view and highlight the connection of v -G-Dedekind domains with their generalization, the (\star, v) -G-Dedekind domains. Now a look at some general properties of DCDs.

It was indicated in [24], using the example of the ring of entire functions, a v -G-Dedekind domain does not behave well under localizations. Using the same example we can conclude that a DCD does not behave well under localization. But let us be clear about that example. Let \mathcal{E} denote the ring of entire functions. Using the fact that \mathcal{E} is a Bezout domain and that an element s of \mathcal{E} that is not divisible by any principal (height one) prime (i.e. one that does not have a zero) is a unit, one can show that $\mathcal{E} = \bigcap_{\mathcal{E}_{(p)}}$ where p ranges over principal height one primes. In fact we have the following lemma.

Lemma 4.2. *Let R be a GCD domain such that an element $s \in R \setminus \{0\}$ is a nonunit if and only if s is divisible by at least one principal prime of R . Then $R = \bigcap_{R_{(p)}}$ where p ranges over principal height one primes of R . Such a ring is completely integrally closed.*

Proof. Obviously if R has no principal height one primes then R is a field and R is trivially an intersection of localizations at its height one primes. So let us assume that R does have a set \wp of principal height one primes and set $S = \bigcap_{p \in \wp} R_{(p)}$. We already have $R \subseteq \bigcap_{R_{(p)}}$. If there is $x \in S \setminus R$, then $x = r/s$ and we can assume that $GCD(r, s) = 1$. Now as $pR_{(p)}$ is of height one and R a GCD domain $R_{(p)}$ is a discrete rank one valuation domain. Next $r/s \in R_{(p)}$ for $p \in \wp$ forces s to be a unit in $R_{(p)}$ meaning s is not divisible by p . But then $x = r/s$ where s is not divisible by any of the $p \in \wp$. This forces s to be a unit, by the rule. But then x is an associate of $r \in R$. Thus $R = \bigcap_{R_{(p)}}$. Finally, each $R_{(p)}$ for each $p \in \wp$ is a discrete rank one valuation domain and hence completely integrally closed and an intersection of completely integrally closed domains is completely integrally closed. \square

Thus we see the reason why one concludes that \mathcal{E} is completely integrally closed. Now \mathcal{E} is Bezout and hence a PVMD such that every nonzero ideal of \mathcal{E} is a t -ideal. Following [17], let P be a nonzero non-maximal prime t -ideal of a PVMD R and set $S = \bigcap_{R_M}$ where M ranges over all the maximal t -ideals which do not contain P . Then $P^{-1} = R_P \cap S$ [17, Proposition 1.1].

Proposition 5. *Let P be a nonzero prime of \mathcal{E} of height greater than one. Then $P_v = \mathcal{E}$.*

Proof. Note that P is contained in no height one prime of \mathcal{E} . But then principal primes are maximal (t -) ideals in \mathcal{E} and so by Lemma 4.2, $S = \mathcal{E}$, forcing $P^{-1} = \mathcal{E}_P \cap \mathcal{E} = \mathcal{E}$. Indeed then $P_v = \mathcal{E}$. Finally if M is a maximal ideal of \mathcal{E} of height greater than one, then M must contain a nonzero non-maximal prime \wp of height greater than one. But then $\mathcal{E} = (\wp)_v \subseteq M_v$. \square

Corollary 7. *In \mathcal{E} it is possible to have a multiplicative set S and an ideal A such that $(A\mathcal{E}_S)_{v'} \subsetneq (A_{v_\mathcal{E}}\mathcal{E}_S)_{v'}$ where v' and $v_\mathcal{E}$ are the v -operations in \mathcal{E}_S and \mathcal{E} respectively.*

Proof. Let P be non-zero non-maximal prime ideal of \mathcal{E} of height greater than one contain a maximal ideal M of \mathcal{E} . Then by Proposition 5, $P_{v_\mathcal{E}} = \mathcal{E}$, where $v_\mathcal{E}$ denotes the v -operation on \mathcal{E} . On the other hand in \mathcal{E}_M which is a valuation ring we have $P\mathcal{E}_M$ a non-maximal prime of \mathcal{E}_M and so must be divisorial being the intersection of all the principal ideals containing $P\mathcal{E}_M$. Thus if v' is the v -operation on \mathcal{E}_M , then $(P\mathcal{E}_M)_{v'} = P\mathcal{E}_M \subsetneq \mathcal{E}_M = (P_{v_\mathcal{E}}\mathcal{E}_M)_{v'}$. \square

We now establish that it is the DC property going missing in localization that is responsible for Corollary 7. For this note that a DCD that is also a $*$ -domain is a v -G-Dedekind domain. Now the $*$ -property, as indicated in [25], fares nicely under localization and so if the v -G-Dedekind property goes missing in localizing, it is the DC property that goes missing in localizing. However, the situation can be brought under control, once we introduce some restriction.

Proposition 6. *If R is DC such that for each ideal $A \in F(R)$ there is a finitely generated $B \subseteq A$ with $A_v = B_v$, then for each multiplicative set S the ring R_S is DC.*

Proof. Note that a domain with the given property is DC because in it for every ideal A we have A_v of finite type. On the other hand a domain with the given property is known to be a Mori domain ([26] and references there) and a Mori domain stays a Mori domain under localization. We include the proof below.

Let S be a multiplicative set in R such that for each $A \in F(R)$ there is a finitely generated $B \subseteq A$ with $A_v = B_v$ and let α be an ideal of R_S . Then $\alpha = AR_S$ where $A = \alpha \cap R$. Let $B \subseteq A$ where B is finitely generated such that $A_v = B_v$. Note that $A \subseteq B_v$ and so $\alpha = AR_S \subseteq B_v R_S$. Thus $\alpha_v = (AR_S)_v \subseteq (B_v R_S)_v = (BR_S)_v$, since B is finitely generated [23]. Since $B \subseteq A$, we have $BR_S \subseteq AR_S$ and so $(BR_S)_v \subseteq (AR_S)_v$. Thus $(BR_S)_v = (AR_S)_v$ with BR_S finitely generated and contained in AR_S . \square

Because in a DCD R , A_v is a finite intersection of principal fractional ideals we conclude that $A_v R_S$ is divisorial. Thus $(AR_S)_v \subseteq A_v R_S$. However as Corollary 7 indicates the inclusion may be strict on occasion. Now generally if $A \in F(R)$ is finitely generated and S a multiplicative set of R , then $(AR_S)_{v'} = (A_{v_R} R_S)_{v'}$, where v' and v_R are v -operations on R_S and R respectively [23]. This is a general formula and so it works for DCDs too, but in a slightly modified form.

Proposition 7. *If $A \in F(R)$ is nonzero finitely generated, S a multiplicative set of R and if R is DC, then $(AR_S)_{v'} = A_{v_R} R_S$, where v' and v_R are v -operations on R_S and R respectively.*

Proof. The proof follows from the fact that A_{v_R} is a finite intertsection of principal fractional ideals and so $A_{v_R}R_S$ is a divisorial ideal of R_S and that makes $A_{v_R}R_S = (A_{v_R}R_S)_{v'}$. \square

A prime example of a DCD is a Mori domain and Roitman [22] has produced an example of a Mori domain R such that $R[X]$ is not Mori. On the other hand for integrally closed integral domains Querre [21] proved the following result.

Theorem 4.3. *An integral domain R is integrally closed if and only if for every integral ideal B of $R[X]$ with $B \cap R \neq 0$, we have $B_v = (A_B[X])_v = (A_B)_v[X]$ where A_B is the ideal of R generated by the coefficients of elements of B .*

For a clearer treatment of Theorem 4.3, see section 3 of [6]. For now, we use this theorem to prove the following result.

Theorem 4.4. *Let R be an integrally closed integral domain. The polynomial ring $R[X]$ is DC if and only if R is.*

Proof. Suppose that R is DC. Then for every ideal $I \in F(R)$ of R we have $I_v = J_v$ where J is finitely generated. Now, let $H \in F(R[X])$. Because R is integrally closed, according to Theorem 2.1 of [6], $H = \frac{f(X)}{g(X)}B$ where $f(X), g(X) \in R[X]$ and B is an ideal of $R[X]$ with $B \cap R \neq (0)$. But then $H_v = \frac{f(X)}{g(X)}B_v = \frac{f(X)}{g(X)}((A_B)_v[X])$ is a v -ideal of finite type because, as A_B is an integral ideal of R , $(A_B)_v$ is a v -ideal of finite type. Conversely let I be a nonzero integral ideal of R and suppose that $R[X]$ is DC. Then $(I[X])_v = I_v[X]$ is a v -ideal of finite type and this forces I_v to be a v -ideal of finite type. \square

REFERENCES

- [1] D.D. Anderson, On the ideal equation $I(B \cap C) = IB \cap IC$, *Canad. Math. Bull.* 26 (1983), 331-332.
- [2] D.D. Anderson and D.F. Anderson, Generalized GCD domains, *Comment. Math. Univ. St. Pauli* 28(1979) 215-221.
- [3] D.D. Anderson, D.F. Anderson, M. Fontana and M. Zafrullah, On v -domains and star operations, *Comm. Algebra* 37 (2009), 3018-3043.
- [4] D.D. Anderson and S. Cook, Two star operations and their induced lattices, *Comm. Algebra*, 28(5) (2000), 2461-2475.
- [5] D.D. Anderson and B.G. Kang, Pseudo-Dedekind domains and divisorial ideals in $R[X]_T$. *J. Algebra* 122 (1989) 323-336.
- [6] D.D. Anderson, D.J. Kwak and M. Zafrullah, Agreeable domains, *Comm. Algebra* 23 (13) (2010), 4861- 4883.
- [7] D.F. Anderson and D. Dobbs, On the product of ideals, *Canad. Math. Bull.* 26(1983) 106-116.
- [8] D.F. Anderson, Integral v -ideals, *Glasgow Math. J.* 22 (1981) 167-172.
- [9] E. Akalan, On generalized Dedekind prime rings, *J. Algebra* 320 (2008) 2907-2916.
- [10] P.M. Cohn, Bézout rings and their subrings, *Proc. Camb. Philos. Soc.* 64 (1968) 251-264.
- [11] J. Elliott, *Rings, modules and closure operations*, Springer Monographs in Mathematics. Springer, Cham, 2019.
- [12] R. Fossum, *The divisor class group of a Krull domain*, *Ergebnisse der Mathematik und ihrer grenzgebiete B.* 74, Springer-Verlag, Berlin, Heidelberg, New York, 1973.
- [13] M. Fontana and S. Gabelli, On the class group and the local class group of a pullback, *J. Algebra* 181 (1996), 803-835.
- [14] M. Fontana, P. Jara and E. Santos, Prufer \ast -star multiplication domains and semistar operations, *J. Algebra Appl.* 2 (2003) 21-50.
- [15] R. Gilmer, *Multiplicative Ideal Theory*. New York: Marcel Dekker (1972).
- [16] F.Halter-Koch, *Ideal Systems, An introduction to ideal theory*, Marcel Dekker, New York, 1998.

- [17] E. Houston, On divisorial prime ideals in Prüfer v -multiplication domains, J. Pure Appl. Algebra 42 (1986) 55-62.
- [18] E. Houston, S. Malik and J. Mott, Characterizations of \star -multiplication domains, Canad. Math. Bull. 27(1)(1984) 48-52.
- [19] P. Jaffard, Les systemes d'ideaux , Dunod, Paris, 1960.
- [20] N. Popescu, On a class of Prüfer domains. Rev. Roumaine Math. Pure Appl. 29 (1984) 777-786.
- [21] J. Querre, Ideaux divisoriels d'un anneau de polynomes, J. Algebra 64 (1980) 270-284.
- [22] M. Roitman, On polynomial extensions of Mori domains over countable fields, J. Pure Appl. Algebra 64 (1990) 315-328.
- [23] M. Zafrullah, Finite conductor domains, Manuscripta Math. 24(1978) 191-203
- [24] M. Zafrullah, (1986). On generalized Dedekind domains. Mathematika 33:285-295.
- [25] M. Zafrullah, On a property of pre-Schreier domains, Comm. Alg. 15 (1987) 1895-1920.
- [26] M. Zafrullah, Ascending chain condition and star operations. Comm. Algebra 17 (1989) 1523-1533.
- [27] M. Zafrullah, Putting t -invertibility to use, Chapter 20, in Non-Noetherian Commutative Ring Theory, pp 429-457, Math. Appl., vol. 520, Kluwer Acad. Publ., Dordrecht, 2000, Editors: S. Glaz and S. Chapman.

DEPARTMENT OF MATHEMATICS, IDAHO STATE UNIVERSITY, POCA TELLO, 83209 ID
E-mail address: `mzafrullah@usa.net`